

Table 1C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Demographic Indicators (2)

Page: 1
Run Date: 7 / 2 / 12

Total Records	Number	%	Source of Data	Number	%
Complete Records	24,890	75.2	WIC	33,109	100.0
Prenatal Records	2,211	6.7	Maternal Health	0	0.0
Postpartum Records	6,008	18.1	Other	0	0.0
Total Records Accepted	33,109		Total	33,109	100.0
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%	Age Distribution	Number	%	Education Distribution	Number	%
White, Not Hispanic	17,639	53.3	< 15 Years	81	0.2	< High School	8,710	28.5
Black, Not Hispanic	6,306	19.0	15 - 17 Years	1,472	4.4	High School	13,813	45.1
Hispanic	6,151	18.6	18 - 19 Years	3,786	11.4	> High School	8,092	26.4
American Indian/Alaska Native	563	1.7	20 - 29 Years	20,521	62.0	Total	30,615	100.0
Asian/Pacific Islander	1,670	5.0	30 - 39 Years	6,795	20.5	<i>Excluded Unknown: 2,494</i>	<i>7.5</i>	
Multiple Races	780	2.4	>= 40 Years	447	1.4			
All Other	0	0.0	Total	33,102	100.0			
Total	33,109	100.0	<i>Excluded Unknown: 7</i>	<i>0.0</i>				
<i>Excluded Unknown: 0</i>	<i>0.0</i>							

% Poverty Level	Number	%	Program Participation (3)	Number	%	Migrant Status	Number	%
0 - 50	6,922	25.6	WIC	27,101	100.0	Yes	41	0.1
51 - 100	5,391	20.0	SNAP	26,737	53.8	No	33,068	99.9
101 - 130	2,694	10.0	Medicaid	27,003	35.8	Total	33,109	100.0
131 - 150	1,358	5.0	TANF	27,003	1.5	<i>Excluded Unknown: 0</i>	<i>0.0</i>	
151 - 185	1,757	6.5	<i>Excluded Unknown:</i>					
186 - 200	0	0.0	WIC:	0	0.0			
> 200	0	0.0	SNAP:	364	1.3			
Adjunctive Eligibility	8,881	32.9	Medicaid:	98	0.4			
Total	27,003	100.0	TANF:	98	0.4			
<i>Excluded Unknown: 98</i>	<i>0.4</i>							

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2011 Pregnancy Nutrition Surveillance
Wisconsin
Summary of Health Indicators

(1)

Page: 2
Run Date: 7 / 2 / 12

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	32,735	3.6	1,106,370	4.5
Normal Weight	32,735	39.5	1,106,370	42.1
Overweight	32,735	25.8	1,106,370	26.0
Obese	32,735	31.0	1,106,370	27.4
Maternal Weight Gain (4)				
Less Than Ideal	26,133	22.2	835,927	21.5
Ideal	26,133	28.1	835,927	30.6
Greater Than Ideal	26,133	49.7	835,927	48.0
Anemia (Low Hb/Hct) (5)				
1st Trimester	10,839	6.4	419,016	7.6
2nd Trimester	8,974	10.6	333,582	12.0
3rd Trimester	4,870	30.8	164,201	33.9
Postpartum	26,649	22.9	346,981	28.1
Parity (6)				
0	31,185	38.5	965,206	40.3
1	31,185	28.2	965,206	26.8
2	31,185	17.5	965,206	16.7
≥ 3	31,185	15.8	965,206	16.3
Interpregnancy Interval				
< 6 Months	16,134	9.8	554,361	11.7
6 - <18 Months	16,134	26.8	554,361	25.9
≥18 Months	16,134	63.4	554,361	62.4
Gestational Diabetes	29,934	5.5	419,956	5.2
Hypertension During Pregnancy	29,934	5.0	467,675	6.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Health Indicators

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	30,406	88.3	429,770	83.1
No Care	30,406	2.3	429,770	4.1
WIC Enrollment				
1st Trimester	33,014	36.0	1,074,863	36.0
2nd Trimester	33,014	29.9	1,074,863	33.9
3rd Trimester	33,014	16.1	1,074,863	16.9
Postpartum (3)	33,014	18.0	1,074,863	13.2
Prepregnancy Multivitamin Consumption (4)	32,041	18.6	380,154	19.2
Multivitamin Consumption During Pregnancy (5)	26,132	86.2	418,448	85.0
Smoking				
3 Months Prior to Pregnancy	31,366	34.3	814,631	23.0
During Pregnancy	27,014	22.2	710,227	14.0
Last 3 Months of Pregnancy	29,887	18.3	715,034	12.1
Postpartum	29,893	19.6	722,233	14.1
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	9,230	36.4	166,084	43.5
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	8,192	32.1	137,883	38.1
Smoking in Household				
During Pregnancy	26,475	16.1	660,570	16.3
Postpartum	29,899	4.9	562,126	15.6
Drinking				
3 Months Prior to Pregnancy	31,250	11.8	930,233	8.8
Last 3 Months of Pregnancy	29,869	0.2	577,187	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Health Indicators

Page: 4
Run Date: 7 / 2 / 12

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	28,973	1.1	880,933	1.1
Low	28,973	6.8	880,933	6.9
Normal	28,973	84.7	880,933	85.3
High	28,973	7.4	880,933	6.7
Preterm (4)	29,995	9.3	948,286	10.7
Full term Low Birth Weight (5)	26,277	3.2	749,107	3.4
Multiple Births	30,746	1.3	978,390	1.4
Breastfeeding				
Ever Breastfed	29,477	72.3	871,811	68.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.



Table 3C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Racial and Ethnic Distribution by Contributor to Nation

Page: 5
Run Date: 7 / 2 / 12

Comparison of Racial and Ethnic Distribution (2)								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wisconsin	33,109	53.3	19.0	18.6	1.7	5.0	2.4	0.0
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Age Distribution by Contributor to Nation

Page: 6
Run Date: 7 / 2 / 12

Comparison of Age Distribution (2)								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wisconsin	33,109	0.2	4.4	11.4	62.0	20.5	1.4	0.0
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Maternal Health Indicators by Contributor to Nation

Page: 7
Run Date: 7 / 2 / 12

Comparison of Maternal Health Indicators (2)														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Wisconsin	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9	29,934	5.5	29,934	5.0
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Maternal Behavioral Indicators by Contributor to Nation

Page: 8
Run Date: 7 / 2 / 12

Comparison of Maternal Behavioral Indicators (2)							
	Medical Care			WIC Enrollment		Multivitamin Consumption	
	1st Trimester		No Care	1st Trimester		Prepregnancy (3)	During Pregnancy (4)
	Number	%	%	Number	%	Number	%
Wisconsin	30,406	88.3	2.3	33,014	36.0	32,041	18.6
Nation (Prior Year)	429,770	83.1	4.1	1,074,863	36.0	380,154	19.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Contributor to Nation

Page: 9
Run Date: 7 / 2 / 12

Comparison of Smoking Indicators (2)										
	Smoking		Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household						
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy		Prenatal	Postpartum					
	Number	%	Number	%	Number	%				
Wisconsin	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Infant Health Indicators by Contributor to Nation

Page: 10
Run Date: 7 / 2 / 12

Comparison of Infant Health Indicators (2)								
	Birthweight			Preterm (5)	Full Term LBW (6)	Ever Breastfed		
	Low (3)		High (4)					
	Number	%	%	Number	%	Number	%	
Wisconsin	28,973	7.9	7.4	29,995	9.3	26,277	3.2	29,477 72.3
Nation (Prior Year)	880,933	8.0	6.7	948,286	10.7	749,107	3.4	871,811 68.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 9C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 11
Run Date: 7 / 2 / 12

Maternal Health Indicators by Race/Ethnicity, Age, or Education

Maternal Health Indicators by Race/Ethnicity, Age, or Education												(2)		
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity														
White, Not Hispanic	17,475	4.0	54.6	13,707	19.7	52.9	2,621	23.3	14,019	15.9	15,929	5.0	15,929	5.1
Black, Not Hispanic	6,205	3.4	61.4	5,057	24.0	49.9	1,037	52.7	5,140	41.1	5,661	4.1	5,661	7.3
Hispanic	6,061	2.7	62.3	5,105	24.8	44.9	713	26.8	5,065	21.2	5,563	8.1	5,563	3.6
American Indian/Alaska Native	560	2.0	68.6	416	23.1	51.4	91	*	398	26.6	506	5.7	506	4.0
Asian/Pacific Islander	1,658	5.1	40.3	1,216	31.8	30.2	298	31.9	1,410	30.9	1,563	6.5	1,563	1.3
Multiple Races	776	3.1	55.7	632	21.0	52.2	110	25.5	617	22.9	712	4.1	712	4.8
All Other/Unknown	0	*	*	0	*	*	0	*	0	*	0	*	0	*
Total	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9	29,934	5.5	29,934	5.0
Age														
< 15 Years	80	*	*	68	*	*	15	*	67	*	74	*	74	*
15 - 17 Years	1,465	8.7	34.2	1,179	19.5	53.6	221	39.8	1,211	32.7	1,355	1.0	1,355	4.0
18 - 19 Years	3,739	5.4	41.2	3,019	19.4	54.1	539	37.3	3,017	27.3	3,417	1.8	3,417	4.8
20 - 29 Years	20,288	3.6	56.9	16,230	22.0	49.5	3,149	30.6	16,558	22.4	18,596	4.4	18,596	4.3
30 - 39 Years	6,710	1.7	69.5	5,307	24.4	47.2	888	26.2	5,469	19.7	6,122	11.2	6,122	6.9
>= 40 Years	446	1.1	74.2	323	30.7	39.6	58	*	327	18.0	363	17.6	363	14.6
Unknown	7	*	*	7	*	*	0	*	0	*	7	*	7	*
Total	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9	29,934	5.5	29,934	5.0
Education														
< High School	8,641	4.3	54.7	7,065	25.4	46.3	1,256	38.5	7,082	27.3	7,899	5.3	7,899	4.3
High School	13,687	3.7	57.3	11,084	21.9	50.4	2,049	29.6	11,115	22.8	12,579	5.0	12,579	5.1
> High School	8,012	2.8	59.3	6,332	18.3	53.0	1,195	25.6	6,540	18.4	7,299	6.4	7,299	6.0
Unknown	2,395	3.5	54.2	1,652	25.1	46.7	370	27.8	1,912	22.1	2,157	6.4	2,157	4.2
Total	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9	29,934	5.5	29,934	5.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 10C
2011 Pregnancy Nutrition Surveillance
Wisconsin
Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education

(1)

Page: 12
Run Date: 7 / 2 / 12

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education ⁽²⁾									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	16,190	91.8	2.0	17,591	35.8	17,155	23.3	13,695	90.4
Black, Not Hispanic	5,749	85.4	1.5	6,296	31.4	6,105	9.9	4,845	83.4
Hispanic	5,672	84.5	4.0	6,126	43.1	5,927	17.2	5,088	81.5
American Indian/Alaska Native	507	84.0	2.0	559	39.7	553	17.4	477	91.0
Asian/Pacific Islander	1,574	79.0	2.4	1,665	26.7	1,539	9.0	1,411	69.2
Multiple Races	714	88.1	1.1	777	37.6	762	13.4	616	88.1
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	30,406	88.3	2.3	33,014	36.0	32,041	18.6	26,132	86.2
Age									
< 15 Years	75	*	*	81	*	79	*	69	*
15 - 17 Years	1,378	77.8	2.5	1,471	36.2	1,443	6.2	1,272	84.7
18 - 19 Years	3,453	85.8	2.5	3,771	41.7	3,679	8.6	3,254	84.3
20 - 29 Years	18,855	89.4	2.1	20,460	34.8	19,838	18.1	16,014	86.4
30 - 39 Years	6,254	89.6	2.5	6,779	36.6	6,569	27.9	5,174	87.4
>= 40 Years	384	83.6	3.4	445	36.4	426	29.3	344	84.0
Unknown	7	*	*	7	*	7	*	5	*
Total	30,406	88.3	2.3	33,014	36.0	32,041	18.6	26,132	86.2
Education									
< High School	8,035	83.4	2.9	8,686	38.1	8,453	11.4	7,181	80.9
High School	12,752	89.3	2.2	13,778	38.0	13,481	16.9	11,171	87.2
> High School	7,437	92.5	1.6	8,071	33.2	7,890	29.1	6,143	91.7
Unknown	2,182	86.8	2.2	2,479	26.8	2,217	19.2	1,637	81.6
Total	30,406	88.3	2.3	33,014	36.0	32,041	18.6	26,132	86.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 11C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Smoking Indicators by Race/Ethnicity, Age, or Education

Page: 13
Run Date: 7 / 2 / 12

Smoking Indicators by Race/Ethnicity, Age, or Education (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%		Number	%	Number	%	
Race/Ethnicity										
White, Not Hispanic	16,894	46.0	15,908	26.0	5,906	29.3	13,914	18.7	15,910	4.9
Black, Not Hispanic	5,824	22.8	5,651	11.9	977	34.9	4,878	20.0	5,658	8.3
Hispanic	5,768	14.1	5,555	5.2	638	49.5	5,156	6.5	5,558	2.1
American Indian/Alaska Native	540	59.6	501	31.3	254	31.5	478	26.6	501	7.6
Asian/Pacific Islander	1,596	11.3	1,562	4.2	154	46.1	1,419	3.6	1,562	2.2
Multiple Races	744	45.4	710	23.4	263	34.2	630	25.6	710	6.3
All Other/Unknown	0	*	0	*	0	*	0	*	0	*
Total	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9
Age										
< 15 Years	76	*	74	*	5	*	70	*	74	*
15 - 17 Years	1,398	24.6	1,353	10.4	286	41.6	1,282	29.1	1,353	11.1
18 - 19 Years	3,628	39.1	3,416	17.1	1,157	37.9	3,300	26.9	3,415	7.7
20 - 29 Years	19,352	36.6	18,563	19.7	5,416	31.9	16,202	14.7	18,572	4.5
30 - 39 Years	6,476	27.9	6,111	17.2	1,271	26.0	5,271	10.3	6,115	3.5
>= 40 Years	430	19.5	363	14.9	57	*	345	12.2	363	3.6
Unknown	6	*	7	*	0	*	5	*	7	*
Total	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9
Education										
< High School	8,189	35.5	7,889	20.7	2,257	28.0	7,252	19.5	7,891	7.5
High School	13,184	38.8	12,563	21.0	3,992	31.4	11,334	17.6	12,570	4.9
> High School	7,734	27.0	7,292	12.3	1,556	39.2	6,228	10.4	7,294	2.8
Unknown	2,259	28.1	2,143	14.9	387	33.9	1,661	11.4	2,144	3.3
Total	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 12C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Infant Health Indicators by Race/Ethnicity, Age, or Education

Infant Health Indicators by Race/Ethnicity, Age, or Education (2)									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	15,404	7.0	8.4	15,947	9.0	14,001	2.6	15,680	74.3
Black, Not Hispanic	5,473	12.5	3.7	5,693	11.8	4,844	5.4	5,552	58.9
Hispanic	5,415	6.3	8.7	5,590	8.4	4,962	2.6	5,531	85.5
American Indian/Alaska Native	476	4.8	13.7	499	7.2	440	1.6	488	69.3
Asian/Pacific Islander	1,518	6.5	3.8	1,558	7.1	1,408	3.1	1,529	54.0
Multiple Races	687	8.2	7.9	708	9.5	622	4.0	697	72.0
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	28,973	7.9	7.4	29,995	9.3	26,277	3.2	29,477	72.3
Age									
< 15 Years	71	*	*	75	*	61	*	73	*
15 - 17 Years	1,294	7.3	3.2	1,354	8.6	1,185	3.8	1,316	66.3
18 - 19 Years	3,323	6.9	5.7	3,426	8.7	3,034	3.6	3,371	70.8
20 - 29 Years	18,043	6.5	7.1	18,665	8.9	16,437	3.0	18,342	72.3
30 - 39 Years	5,889	7.3	10.2	6,110	10.7	5,260	3.3	6,018	74.8
>= 40 Years	346	13.0	9.0	358	15.1	293	4.8	350	69.7
Unknown	7	*	*	7	*	7	*	7	*
Total	28,973	7.9	7.4	29,995	9.3	26,277	3.2	29,477	72.3
Education									
< High School	7,623	7.4	6.4	7,902	9.3	6,923	3.8	7,740	67.0
High School	12,175	6.6	7.1	12,569	9.1	11,078	3.1	12,339	70.1
> High School	7,078	6.3	8.9	7,305	9.6	6,395	2.6	7,228	81.7
Unknown	2,097	7.4	8.0	2,219	9.8	1,881	3.6	2,170	72.5
Total	28,973	7.9	7.4	29,995	9.3	26,277	3.2	29,477	72.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 13C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Maternal Weight Gain and Birth Outcomes by Select Health Indicators

Page: 15
Run Date: 7 / 2 / 12

Maternal Weight Gain and Birth Outcomes by Select Health Indicators (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal	> Ideal		Birthweight			Preterm (6)		Full Term LBW (7)	
				Low (4)		High (5)				
	Number	%	%	Number	%	%	Number	%	Number	%
Prepregnancy BMI										
Underweight	954	31.1	27.9	1,015	12.3	2.1	1,060	11.8	899	5.9
Normal Weight	10,377	24.9	40.7	11,363	8.5	4.8	11,738	9.1	10,329	3.6
Overweight	6,789	14.6	60.7	7,457	7.5	7.8	7,689	9.0	6,785	2.8
Obese	8,013	24.0	54.4	8,921	6.9	10.8	9,189	9.5	8,073	2.6
Weight Gain										
< Ideal	-	-	-	5,252	13.2	3.9	5,439	13.2	4,561	5.9
Ideal	-	-	-	7,096	8.2	5.2	7,300	9.3	6,437	3.4
> Ideal	-	-	-	12,517	5.3	9.9	12,895	7.3	11,594	2.0
Smoking During Pregnancy										
Smokers	4,554	24.1	49.8	5,102	10.0	5.2	5,260	9.4	4,632	5.2
Non-Smokers	16,668	21.6	49.7	18,441	6.5	8.2	18,894	8.3	16,924	2.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Racial and Ethnic Distribution

Page: 16
Run Date: 7 / 2 / 12

Summary of Trends in Racial and Ethnic Distribution (2)								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2011	33,109	53.3	19.0	18.6	1.7	5.0	2.4	0.0
2010	33,789	53.4	19.0	18.9	1.8	4.7	2.2	0.0
2009	35,896	53.7	19.1	18.9	1.9	4.5	1.9	0.0
2008	35,549	53.7	19.1	19.2	1.7	4.5	1.8	0.0
2007	34,934	53.6	19.3	19.1	2.0	4.5	1.7	0.0
2006	31,077	52.1	18.9	19.5	2.0	4.2	1.6	1.7
2005	29,354	54.7	19.2	17.2	2.2	4.3	0.4	2.0
2004	33,697	57.6	19.4	16.4	2.2	4.2	0.0	0.2
2003	0	*	*	*	*	*	*	*
2002	28,924	58.6	19.7	15.7	2.2	3.8	0.0	0.0
2001	31,080	58.3	20.0	15.5	2.2	4.0	0.0	0.0
2000	38,155	59.3	20.8	13.6	2.1	4.2	0.0	0.0
1999	30,872	61.1	20.9	11.9	2.3	3.9	0.0	0.0
1998	30,319	61.9	21.4	10.5	2.1	4.1	0.0	0.0
1997	30,380	62.3	21.4	9.5	2.4	4.3	0.0	0.0
1996	12,251	84.5	4.2	5.3	1.0	5.1	0.0	0.0
1995	29,036	62.4	21.8	9.0	2.1	4.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 15C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution (2)								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2011	33,109	0.2	4.4	11.4	62.0	20.5	1.4	0.0
2010	33,789	0.2	5.1	12.5	61.5	19.4	1.3	0.0
2009	35,896	0.3	5.5	13.0	61.8	18.3	1.2	0.0
2008	35,549	0.3	6.2	13.4	61.6	17.2	1.1	0.0
2007	34,934	0.3	6.6	13.8	61.8	16.2	1.2	0.1
2006	31,077	0.4	6.7	14.1	62.1	15.5	1.1	0.1
2005	29,354	0.5	6.7	14.0	62.0	15.7	1.2	0.0
2004	33,697	0.5	6.7	14.7	61.0	16.0	1.2	0.0
2003	0	*	*	*	*	*	*	*
2002	28,924	0.5	7.4	15.9	59.6	15.5	1.1	0.0
2001	31,080	0.3	6.6	14.6	60.6	16.8	1.2	0.0
2000	38,155	0.5	8.0	16.4	58.7	15.3	1.1	0.0
1999	30,872	0.6	8.6	16.8	57.7	15.3	1.1	0.0
1998	30,319	0.7	8.8	16.6	57.2	15.6	1.0	0.0
1997	30,380	0.8	9.5	16.0	56.8	15.9	1.0	0.0
1996	12,251	0.4	7.2	15.7	58.7	17.0	1.0	0.0
1995	29,036	1.0	9.7	15.9	56.2	16.1	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 16C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators (2)										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2011	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9
2010	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9
2009	35,482	3.7	55.1	26,554	21.6	50.9	5,697	31.9	28,469	25.0
2008	35,138	3.7	53.9	24,071	20.6	51.5	5,983	32.5	29,357	24.8
2007	34,503	3.9	53.1	24,133	21.0	50.7	5,810	33.1	29,138	25.8
2006	30,733	4.4	53.0	21,617	21.9	50.5	4,588	29.4	24,349	24.9
2005	25,573	4.5	50.4	21,542	24.5	45.5	4,675	30.4	19,010	25.3
2004	29,546	4.7	50.1	26,190	23.9	45.9	5,082	33.2	16,237	26.1
2003	0	*	*	0	*	*	0	*	0	*
2002	25,042	5.1	48.2	21,724	23.0	46.1	4,619	31.0	13,207	25.7
2001	27,218	5.4	47.2	23,533	23.7	45.1	2,855	17.0	14,327	26.4
2000	32,882	5.8	45.8	28,239	24.6	44.8	5,363	26.1	14,148	26.4
1999	27,032	6.0	44.9	23,018	25.2	44.0	4,840	24.5	10,218	25.6
1998	26,849	5.7	44.6	23,617	24.4	43.7	4,515	29.7	9,529	23.2
1997	26,105	5.8	43.7	23,105	25.0	43.1	4,565	27.7	9,599	21.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 16C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Maternal Health Indicators

Page: 19
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Health Indicators (2)										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1996	10,183	6.3	41.5	9,004	25.7	41.7	1,775	22.3	3,636	18.2
1995	24,836	6.4	40.5	21,703	26.0	42.0	4,377	31.9	8,906	21.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2011	30,406	88.3	2.3	33,014	36.0
2010	31,228	87.2	2.5	33,671	35.8
2009	33,210	85.6	2.8	35,778	34.1
2008	32,437	84.0	2.3	34,433	33.0
2007	31,285	83.3	2.3	33,775	33.7
2006	11,903	83.9	0.1	27,099	29.8
2005	22,622	74.9	12.5	28,229	33.2
2004	30,590	74.3	13.9	32,024	34.7
2003	0	*	*	0	*
2002	25,779	74.1	12.5	27,533	33.2
2001	27,652	74.3	10.7	30,274	0.0
2000	33,694	72.5	12.4	36,735	21.1
1999	26,697	71.1	13.0	29,540	23.2
1998	26,953	71.1	12.4	28,767	36.1
1997	27,168	71.3	12.2	28,118	34.4
1996	10,761	72.9	12.9	11,274	36.0
1995	25,376	70.1	13.3	25,878	32.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 18C
2011 Pregnancy Nutrition Surveillance
Wisconsin
Summary of Trends in Smoking Indicators

(1)

Page: 21
Run Date: 7 / 2 / 12

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2011	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9
2010	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
2009	34,096	36.1	32,593	19.7	9,317	31.5	28,409	20.4	32,610	6.9
2008	33,969	36.6	32,190	20.4	9,493	30.9	28,619	21.6	32,204	7.9
2007	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
2006	20,837	33.1	28,206	21.2	6,146	26.5	14,138	25.1	20,724	11.0
2005	18,399	39.8	22,131	26.4	6,285	30.0	18,584	39.8	26,480	33.6
2004	26,351	40.8	29,526	23.7	9,094	29.3	26,414	41.6	30,700	39.6
2003	0	*	0	*	0	*	0	*	0	*
2002	22,548	42.2	24,910	24.5	6,872	32.9	22,642	43.4	25,745	41.5
2001	23,803	42.5	26,637	25.5	8,350	26.8	23,769	44.9	27,572	42.9
2000	30,624	42.8	31,613	25.6	10,461	27.6	30,764	46.6	33,784	44.9
1999	24,100	44.4	25,437	26.2	8,424	28.5	24,204	47.1	27,080	45.2
1998	29,369	43.2	6,139	25.4	212	26.9	24,115	47.0	6,326	42.3
1997	29,464	42.2	6,059	25.4	0	*	23,678	47.4	6,332	40.4
1996	11,796	45.3	2,505	27.3	0	*	9,450	51.1	2,622	43.1
1995	27,869	41.5	6,453	26.2	0	*	21,792	49.0	6,828	43.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 19C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2011	28,973	7.9	7.4	29,995	9.3	26,277	3.2	29,477	72.3
2010	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
2009	31,539	7.8	7.8	32,660	9.7	28,463	3.1	32,009	69.0
2008	31,033	7.7	7.8	31,344	10.2	27,048	3.1	31,499	67.0
2007	30,255	7.8	7.8	30,697	10.4	26,286	3.2	30,747	65.8
2006	26,732	7.7	7.7	23,522	10.8	20,880	3.0	27,572	64.1
2005	24,726	7.8	8.2	24,324	11.1	21,339	2.9	24,996	62.3
2004	28,209	7.6	8.6	27,611	11.4	24,125	3.0	28,189	62.2
2003	0	*	*	0	*	0	*	0	*
2002	23,758	7.4	9.2	23,341	10.7	20,350	3.1	23,977	59.2
2001	25,501	7.4	9.2	12,367	27.9	8,615	8.5	25,708	57.2
2000	29,964	7.3	9.4	23,960	12.8	20,250	3.9	30,279	55.3
1999	24,383	7.5	9.4	20,706	15.5	17,201	3.6	23,396	51.1
1998	25,133	7.6	9.3	24,536	11.6	21,191	2.8	20,339	42.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Infant Health Indicators

Page: 23
Run Date: 7 / 2 / 12

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1997	24,451	7.6	9.5	24,093	11.2	20,374	3.1	21,352	43.2
1996	9,702	6.4	11.0	9,469	10.3	8,208	2.5	9,966	56.0
1995	23,260	7.3	9.6	21,362	11.2	18,699	3.1	23,216	50.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 24
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2011	17,475	4.0	54.6	13,707	19.7	52.9	2,621	23.3	14,019	15.9
	2010	17,871	3.9	54.3	14,056	19.3	53.1	2,867	23.5	14,248	16.7
	2009	19,076	4.3	53.0	14,596	18.4	54.9	3,136	25.0	15,122	17.7
	2008	18,891	4.1	51.9	13,316	17.9	55.1	3,223	24.0	15,677	17.5
	2007	18,521	4.6	50.9	13,326	17.9	54.4	3,130	25.4	15,672	18.7
	2006	16,072	5.1	50.2	11,870	18.6	54.4	2,511	22.4	13,019	17.5
	2005	14,082	5.4	47.7	12,245	21.4	48.7	2,574	22.8	10,446	18.5
	2004	17,132	5.4	47.3	15,317	21.3	48.8	2,882	25.7	8,996	18.9
	2003	0	*	*	0	*	*	0	*	0	*
	2002	14,802	5.8	45.6	13,020	21.3	48.6	2,593	22.6	7,489	18.1
Black, Not Hispanic	2011	6,205	3.4	61.4	5,057	24.0	49.9	1,037	52.7	5,140	41.1
	2010	6,334	3.2	61.5	5,169	27.1	47.3	1,150	54.8	5,212	42.4
	2009	6,764	3.0	59.5	4,535	25.1	50.0	1,173	53.3	5,488	43.9
	2008	6,706	3.6	58.7	3,818	23.1	52.1	1,282	53.8	5,556	44.8
	2007	6,649	3.2	59.3	3,986	23.8	51.0	1,252	54.8	5,530	45.1
	2006	5,804	3.8	60.5	3,864	25.7	50.2	902	52.1	4,483	45.2
	2005	5,101	3.4	58.2	4,408	26.9	45.9	965	52.3	3,563	43.4
	2004	5,957	4.0	57.3	5,329	26.9	45.2	1,092	55.3	3,010	46.4
	2003	0	*	*	0	*	*	0	*	0	*
	2002	5,084	4.3	55.7	4,320	24.3	46.3	949	54.9	2,321	44.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 25
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2011	6,061	2.7	62.3	5,105	24.8	44.9	713	26.8	5,065	21.2
	2010	6,292	2.2	60.8	5,226	25.9	43.8	808	31.7	5,267	22.3
	2009	6,688	2.5	58.9	5,336	25.4	44.4	888	28.8	5,472	24.2
	2008	6,696	2.1	57.7	5,076	23.7	45.5	958	31.8	5,725	23.0
	2007	6,528	2.8	56.2	4,961	23.6	44.5	901	29.7	5,616	24.6
	2006	5,978	2.7	56.9	4,202	24.1	44.5	794	26.4	4,916	24.0
	2005	4,217	2.6	53.4	3,407	30.5	37.4	728	26.0	3,610	24.3
	2004	4,532	2.8	53.9	3,903	26.8	40.7	730	31.1	3,117	25.5
	2003	0	*	*	0	*	*	0	*	0	*
	2002	3,658	3.0	50.0	3,088	25.5	40.3	741	29.8	2,650	28.2
American Indian/Alaska Native	2011	560	2.0	68.6	416	23.1	51.4	91	*	398	26.6
	2010	607	2.1	68.0	445	22.7	55.7	104	32.7	409	28.1
	2009	673	1.2	64.8	506	21.3	49.8	91	*	499	26.3
	2008	619	3.9	63.5	450	19.3	51.1	89	*	477	26.6
	2007	683	1.8	60.6	473	21.1	55.0	106	37.7	526	32.5
	2006	619	3.1	62.7	420	22.9	51.2	67	*	466	25.8
	2005	579	2.1	62.0	511	22.1	52.3	85	*	431	28.3
	2004	670	2.8	61.0	582	22.2	51.4	95	*	379	24.8
	2003	0	*	*	0	*	*	0	*	0	*
	2002	575	4.0	59.7	499	20.6	52.7	81	*	210	35.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 26
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2011	1,658	5.1	40.3	1,216	31.8	30.2	298	31.9	1,410	30.9
	2010	1,572	5.3	41.5	1,163	32.5	30.9	268	29.9	1,365	33.4
	2009	1,614	4.8	40.8	1,081	32.8	32.8	308	29.5	1,356	32.7
	2008	1,572	5.8	35.8	970	33.2	28.9	308	34.7	1,387	33.4
	2007	1,543	5.4	36.0	1,012	36.2	28.6	324	31.2	1,329	31.2
	2006	1,309	6.0	34.2	886	36.6	28.3	243	31.3	1,080	32.9
	2005	1,073	6.2	32.5	896	34.9	26.5	260	33.8	858	35.5
	2004	1,203	5.7	33.2	1,058	36.3	24.7	278	31.3	730	33.2
	2003	0	*	*	0	*	*	0	*	0	*
	2002	923	7.5	33.9	797	37.4	23.3	255	29.8	537	37.2
Multiple Races	2011	776	3.1	55.7	632	21.0	52.2	110	25.5	617	22.9
	2010	739	3.7	58.3	597	24.0	50.3	112	28.6	597	25.1
	2009	667	3.3	55.6	500	18.0	53.6	101	31.7	532	24.8
	2008	653	2.9	56.5	441	17.7	57.4	123	35.8	535	23.9
	2007	577	3.3	55.3	375	21.9	53.9	97	*	465	25.4
	2006	486	3.3	50.8	347	21.9	51.9	67	*	385	26.2
	2005	86	*	*	72	*	*	19	*	76	*
	2004	1	*	*	1	*	*	0	*	1	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 27
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2011	0	*	*	0	*	*	0	*	0	*
	2010	0	*	*	0	*	*	0	*	0	*
	2009	0	*	*	0	*	*	0	*	0	*
	2008	1	*	*	0	*	*	0	*	0	*
	2007	2	*	*	0	*	*	0	*	0	*
	2006	465	5.6	48.6	28	*	*	4	*	0	*
	2005	435	6.7	45.7	3	*	*	44	*	26	*
	2004	51	*	*	0	*	*	5	*	4	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
Total All Race/Ethnic Groups	2011	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9
	2010	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9
	2009	35,482	3.7	55.1	26,554	21.6	50.9	5,697	31.9	28,469	25.0
	2008	35,138	3.7	53.9	24,071	20.6	51.5	5,983	32.5	29,357	24.8
	2007	34,503	3.9	53.1	24,133	21.0	50.7	5,810	33.1	29,138	25.8
	2006	30,733	4.4	53.0	21,617	21.9	50.5	4,588	29.4	24,349	24.9
	2005	25,573	4.5	50.4	21,542	24.5	45.5	4,675	30.4	19,010	25.3
	2004	29,546	4.7	50.1	26,190	23.9	45.9	5,082	33.2	16,237	26.1
	2003	0	*	*	0	*	*	0	*	0	*
	2002	25,042	5.1	48.2	21,724	23.0	46.1	4,619	31.0	13,207	25.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 28
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity (2)						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2011	16,190	91.8	2.0	17,591	35.8
	2010	16,610	90.8	2.0	17,992	35.9
	2009	17,837	89.7	2.0	19,219	33.5
	2008	17,417	88.9	1.9	18,530	33.0
	2007	16,803	88.0	1.9	18,195	33.6
	2006	6,548	88.8	0.0	14,333	28.6
	2005	12,797	80.5	10.5	15,420	33.9
	2004	17,664	79.9	11.4	18,387	35.6
	2003	0	*	*	0	*
	2002	15,218	80.2	10.0	16,080	34.7
Black, Not Hispanic	2011	5,749	85.4	1.5	6,296	31.4
	2010	5,923	84.0	1.6	6,405	30.4
	2009	6,324	81.3	2.5	6,834	30.7
	2008	6,203	78.9	2.3	6,490	29.3
	2007	6,015	78.7	2.5	6,409	30.2
	2006	2,401	78.6	0.0	5,031	28.3
	2005	4,516	71.7	10.1	5,478	30.6
	2004	5,927	71.7	11.8	6,346	32.3
	2003	0	*	*	0	*
	2002	4,911	70.2	11.4	5,505	29.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 29
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity (2)						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2011	5,672	84.5	4.0	6,126	43.1
	2010	5,951	82.8	4.5	6,341	43.5
	2009	6,266	81.0	5.1	6,756	41.5
	2008	6,202	79.4	3.3	6,641	38.4
	2007	5,934	79.0	3.0	6,444	39.3
	2006	1,904	81.0	0.2	5,269	33.3
	2005	3,751	63.4	22.7	4,852	35.1
	2004	5,040	61.7	24.8	5,239	37.3
	2003	0	*	*	0	*
	2002	4,096	61.2	22.7	4,303	34.9
American Indian/Alaska Native	2011	507	84.0	2.0	559	39.7
	2010	555	84.1	2.5	607	38.7
	2009	621	82.8	3.2	674	40.1
	2008	550	81.6	3.5	586	36.7
	2007	598	75.8	2.5	656	39.2
	2006	294	70.7	0.0	519	36.8
	2005	520	78.7	4.8	616	42.4
	2004	660	72.7	10.8	695	44.0
	2003	0	*	*	0	*
	2002	550	75.3	8.9	593	40.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 30
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity (2)						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2011	1,574	79.0	2.4	1,665	26.7
	2010	1,503	77.5	3.3	1,581	25.0
	2009	1,530	74.9	3.5	1,623	21.7
	2008	1,472	69.2	2.7	1,542	22.5
	2007	1,411	68.4	3.2	1,502	22.4
	2006	562	66.5	0.0	1,069	18.1
	2005	996	59.0	13.1	1,203	20.2
	2004	1,298	58.3	17.1	1,338	18.7
	2003	0	*	*	0	*
	2002	1,004	53.1	15.0	1,052	18.3
Multiple Races	2011	714	88.1	1.1	777	37.6
	2010	686	86.9	2.0	745	37.6
	2009	632	87.5	1.9	672	36.0
	2008	593	81.8	1.7	644	37.7
	2007	524	81.5	3.4	569	38.0
	2006	194	82.0	0.0	432	27.8
	2005	42	*	*	102	24.5
	2004	1	*	*	0	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 31
Run Date: 7 / 2 / 12

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity (2)						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2011	0	*	*	0	*
	2010	0	*	*	0	*
	2009	0	*	*	0	*
	2008	0	*	*	0	*
	2007	0	*	*	0	*
	2006	0	*	*	446	65.0
	2005	0	*	*	558	44.6
	2004	0	*	*	19	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
Total All Race/Ethnic Groups	2011	30,406	88.3	2.3	33,014	36.0
	2010	31,228	87.2	2.5	33,671	35.8
	2009	33,210	85.6	2.8	35,778	34.1
	2008	32,437	84.0	2.3	34,433	33.0
	2007	31,285	83.3	2.3	33,775	33.7
	2006	11,903	83.9	0.1	27,099	29.8
	2005	22,622	74.9	12.5	28,229	33.2
	2004	30,590	74.3	13.9	32,024	34.7
	2003	0	*	*	0	*
	2002	25,779	74.1	12.5	27,533	33.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 32
Run Date: 7 / 2 / 12

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2011	16,894	46.0	15,908	26.0	5,906	29.3	13,914	18.7	15,910	4.9
	2010	17,309	45.5	16,321	25.7	5,976	29.6	14,096	20.0	16,330	5.3
	2009	18,512	48.4	17,536	27.8	6,773	28.7	14,880	23.7	17,551	7.2
	2008	18,401	49.3	17,301	29.0	6,896	28.2	14,914	25.2	17,312	8.0
	2007	17,872	48.3	16,939	30.7	6,495	25.7	14,581	26.5	16,933	9.2
	2006	10,856	46.6	15,000	30.3	4,548	24.4	7,370	29.6	11,021	11.6
	2005	10,309	54.4	12,936	35.2	4,844	28.5	10,372	49.5	14,813	41.5
	2004	14,915	54.7	17,033	31.6	6,932	28.0	14,929	51.1	17,671	48.3
	2003	0	*	0	*	0	*	0	*	0	*
	2002	12,939	56.8	14,651	32.3	5,350	32.2	12,961	52.8	15,175	49.9
Black, Not Hispanic	2011	5,824	22.8	5,651	11.9	977	34.9	4,878	20.0	5,658	8.3
	2010	5,906	23.8	5,840	12.7	1,087	35.6	5,010	22.1	5,844	8.8
	2009	6,370	24.5	6,197	12.8	1,117	35.3	5,350	26.0	6,197	10.3
	2008	6,417	25.4	6,123	14.0	1,212	33.3	5,518	27.1	6,123	12.5
	2007	6,318	24.2	6,023	13.9	1,116	30.4	5,408	27.1	6,015	13.3
	2006	3,921	22.6	5,379	14.2	745	27.2	2,598	29.4	3,935	16.3
	2005	3,604	25.8	4,190	18.7	762	27.4	3,668	34.3	5,199	30.0
	2004	5,228	27.6	5,669	17.8	1,187	27.0	5,266	36.8	5,954	35.1
	2003	0	*	0	*	0	*	0	*	0	*
	2002	4,477	28.7	4,741	19.1	866	27.4	4,511	38.4	4,922	36.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 33
Run Date: 7 / 2 / 12

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2011	5,768	14.1	5,555	5.2	638	49.5	5,156	6.5	5,558	2.1
	2010	6,018	14.7	5,821	5.0	690	53.5	5,393	7.7	5,820	2.3
	2009	6,362	14.4	6,101	5.5	731	48.4	5,697	8.1	6,102	2.6
	2008	6,386	14.4	6,153	5.1	734	50.3	5,732	9.0	6,152	3.4
	2007	6,227	13.4	6,026	6.1	653	45.3	5,576	10.2	6,011	3.8
	2006	4,269	12.0	5,552	5.3	455	44.0	2,912	12.7	4,051	5.2
	2005	3,228	11.7	3,526	6.1	319	52.7	3,274	17.2	4,610	14.4
	2004	4,483	13.5	4,896	5.7	516	51.2	4,492	18.1	5,077	17.3
	2003	0	*	0	*	0	*	0	*	0	*
2002	3,722	12.1	3,986	4.8	343	56.0	3,742	20.1	4,081	19.3	
American Indian/Alaska Native	2011	540	59.6	501	31.3	254	31.5	478	26.6	501	7.6
	2010	592	61.0	545	33.0	295	34.9	529	27.0	545	6.1
	2009	647	58.4	611	32.9	301	28.9	567	33.9	612	13.1
	2008	601	59.9	545	32.1	278	29.9	529	30.8	547	9.7
	2007	648	55.1	612	32.5	284	26.1	580	32.9	605	11.7
	2006	437	50.8	562	33.8	196	24.5	302	37.4	425	11.3
	2005	453	68.2	509	43.4	271	32.5	455	58.9	578	50.7
	2004	628	65.1	628	41.6	345	26.1	629	61.7	663	58.4
	2003	0	*	0	*	0	*	0	*	0	*
2002	532	64.3	530	40.6	247	30.8	546	59.2	553	57.9	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 34
Run Date: 7 / 2 / 12

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2011	1,596	11.3	1,562	4.2	154	46.1	1,419	3.6	1,562	2.2
	2010	1,515	12.1	1,502	5.3	150	40.0	1,292	5.2	1,502	2.5
	2009	1,564	10.7	1,532	4.0	138	48.6	1,356	4.5	1,532	2.3
	2008	1,528	9.5	1,482	4.6	112	35.7	1,353	6.6	1,483	3.3
	2007	1,467	9.9	1,451	5.2	115	47.8	1,312	7.5	1,448	3.9
	2006	1,002	8.4	1,251	3.8	75	*	714	8.0	945	5.7
	2005	799	12.6	930	6.5	87	*	803	21.0	1,180	17.5
	2004	1,097	12.0	1,299	5.1	114	45.6	1,098	21.1	1,334	20.0
	2003	0	*	0	*	0	*	0	*	0	*
2002	878	9.7	1,002	4.9	66	*	882	21.3	1,014	20.4	
Multiple Races	2011	744	45.4	710	23.4	263	34.2	630	25.6	710	6.3
	2010	708	48.2	678	25.2	270	32.2	611	23.9	679	7.2
	2009	641	49.0	616	25.8	257	35.0	559	28.4	616	9.9
	2008	636	49.8	586	25.3	261	37.5	573	29.1	587	9.9
	2007	561	49.0	524	30.3	219	25.6	498	32.1	522	9.4
	2006	346	41.9	458	25.3	127	26.0	236	27.1	343	16.3
	2005	6	*	40	*	2	*	12	*	100	22.0
	2004	0	*	1	*	0	*	0	*	1	*
	2003	0	*	0	*	0	*	0	*	0	*
2002	0	*	0	*	0	*	0	*	0	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 35
Run Date: 7 / 2 / 12

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2011	0	*	0	*	0	*	0	*	0	*
	2010	0	*	0	*	0	*	0	*	0	*
	2009	0	*	0	*	0	*	0	*	0	*
	2008	0	*	0	*	0	*	0	*	0	*
	2007	1	*	0	*	0	*	1	*	0	*
	2006	6	*	4	*	0	*	6	*	4	*
	2005	0	*	0	*	0	*	0	*	0	*
	2004	0	*	0	*	0	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2011	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9
	2010	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
	2009	34,096	36.1	32,593	19.7	9,317	31.5	28,409	20.4	32,610	6.9
	2008	33,969	36.6	32,190	20.4	9,493	30.9	28,619	21.6	32,204	7.9
	2007	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
	2006	20,837	33.1	28,206	21.2	6,146	26.5	14,138	25.1	20,724	11.0
	2005	18,399	39.8	22,131	26.4	6,285	30.0	18,584	39.8	26,480	33.6
	2004	26,351	40.8	29,526	23.7	9,094	29.3	26,414	41.6	30,700	39.6
	2003	0	*	0	*	0	*	0	*	0	*
	2002	22,548	42.2	24,910	24.5	6,872	32.9	22,642	43.4	25,745	41.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 23C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 36
Run Date: 7 / 2 / 12

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity (2)										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2011	15,404	7.0	8.4	15,947	9.0	14,001	2.6	15,680	74.3
	2010	15,770	7.2	8.3	16,363	9.6	14,251	2.6	16,006	72.0
	2009	16,963	6.9	8.7	17,558	9.1	15,412	2.6	17,202	70.1
	2008	16,660	7.3	9.1	16,874	9.7	14,628	2.9	16,932	67.4
	2007	16,266	7.1	8.9	16,549	10.0	14,278	2.8	16,505	65.9
	2006	14,311	7.3	8.7	12,778	10.0	11,460	3.0	14,678	64.1
	2005	13,898	7.1	9.1	13,584	10.4	12,048	2.5	13,937	62.5
	2004	16,293	7.1	9.5	15,806	10.9	13,884	2.8	16,277	62.9
	2003	0	*	*	0	*	0	*	0	*
	2002	14,008	6.7	10.5	13,660	10.1	12,037	2.8	14,083	60.2
Black, Not Hispanic	2011	5,473	12.5	3.7	5,693	11.8	4,844	5.4	5,552	58.9
	2010	5,598	12.6	3.4	5,846	13.1	4,861	5.2	5,686	57.1
	2009	5,959	12.6	4.1	6,208	13.0	5,175	5.4	6,026	55.6
	2008	5,826	11.5	3.8	5,863	12.4	4,873	5.0	5,890	54.0
	2007	5,650	12.2	4.2	5,752	13.0	4,674	5.5	5,740	52.3
	2006	4,937	11.5	4.2	4,262	14.0	3,638	4.7	5,219	49.7
	2005	4,735	11.9	5.0	4,712	14.4	3,982	4.6	4,839	48.6
	2004	5,441	11.3	5.2	5,469	14.6	4,602	4.9	5,390	46.4
	2003	0	*	*	0	*	0	*	0	*
	2002	4,464	11.4	4.8	4,519	13.3	3,760	4.8	4,579	44.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 37
Run Date: 7 / 2 / 12

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity (2)										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2011	5,415	6.3	8.7	5,590	8.4	4,962	2.6	5,531	85.5
	2010	5,685	5.6	7.7	5,840	8.1	5,202	2.3	5,771	83.6
	2009	5,941	6.0	8.9	6,127	8.3	5,434	2.3	6,077	83.2
	2008	6,015	5.5	8.7	6,066	8.7	5,373	2.0	6,123	83.2
	2007	5,871	5.8	8.8	5,902	9.4	5,170	2.2	6,000	83.0
	2006	5,322	5.7	8.9	4,661	10.3	4,150	1.8	5,453	81.8
	2005	4,299	5.7	9.0	4,217	9.9	3,762	2.5	4,349	81.4
	2004	4,660	5.1	9.3	4,558	9.1	4,098	2.0	4,688	81.4
	2003	0	*	*	0	*	0	*	0	*
	2002	3,817	5.6	9.5	3,726	9.4	3,312	2.7	3,862	77.6
American Indian/Alaska Native	2011	476	4.8	13.7	499	7.2	440	1.6	488	69.3
	2010	530	4.7	12.5	540	8.0	486	1.4	536	66.2
	2009	591	3.4	14.2	612	8.5	541	1.1	596	66.9
	2008	518	5.6	12.7	512	13.1	422	0.9	522	62.1
	2007	572	5.8	14.3	580	11.6	486	3.1	582	59.1
	2006	534	6.2	12.9	452	9.5	406	1.7	549	65.8
	2005	540	6.3	16.5	531	11.5	464	1.9	532	62.0
	2004	588	5.4	14.3	563	13.5	477	1.5	577	60.8
	2003	0	*	*	0	*	0	*	0	*
	2002	509	4.1	15.5	480	11.7	418	1.2	481	61.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin

Page: 38
Run Date: 7 / 2 / 12

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity (2)										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2011	1,518	6.5	3.8	1,558	7.1	1,408	3.1	1,529	54.0
	2010	1,446	7.9	3.0	1,512	7.9	1,335	3.7	1,464	53.8
	2009	1,494	6.4	4.4	1,540	7.5	1,378	3.0	1,506	52.3
	2008	1,445	7.0	3.5	1,456	10.4	1,261	2.8	1,458	47.0
	2007	1,398	6.8	3.5	1,406	8.3	1,237	3.5	1,416	49.6
	2006	1,189	5.9	4.7	982	8.1	896	2.7	1,218	44.8
	2005	1,141	7.2	4.3	1,104	9.6	991	3.2	1,148	45.6
	2004	1,226	7.0	5.6	1,215	9.9	1,064	3.0	1,254	49.5
	2003	0	*	*	0	*	0	*	0	*
	2002	960	7.3	5.1	956	11.9	823	3.9	972	42.2
Multiple Races	2011	687	8.2	7.9	708	9.5	622	4.0	697	72.0
	2010	655	8.2	7.9	683	9.2	597	3.0	668	70.7
	2009	591	10.7	7.6	615	11.5	523	5.0	602	70.3
	2008	569	8.3	8.1	573	12.0	491	2.6	574	67.4
	2007	498	7.8	7.0	508	9.4	441	3.4	504	65.3
	2006	439	10.0	5.9	387	14.5	330	4.2	452	64.6
	2005	95	*	*	93	*	81	*	100	68.0
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Infant Health Indicators by Race/Ethnicity

Page: 39
Run Date: 7 / 2 / 12

Summary of Trends in Infant Health Indicators by Race/Ethnicity										(2)
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2011	0	*	*	0	*	0	*	0	*
	2010	0	*	*	0	*	0	*	0	*
	2009	0	*	*	0	*	0	*	0	*
	2008	0	*	*	0	*	0	*	0	*
	2007	0	*	*	0	*	0	*	0	*
	2006	0	*	*	0	*	0	*	3	*
	2005	18	*	*	83	*	11	*	91	*
	2004	1	*	*	0	*	0	*	3	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2011	28,973	7.9	7.4	29,995	9.3	26,277	3.2	29,477	72.3
	2010	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
	2009	31,539	7.8	7.8	32,660	9.7	28,463	3.1	32,009	69.0
	2008	31,033	7.7	7.8	31,344	10.2	27,048	3.1	31,499	67.0
	2007	30,255	7.8	7.8	30,697	10.4	26,286	3.2	30,747	65.8
	2006	26,732	7.7	7.7	23,522	10.8	20,880	3.0	27,572	64.1
	2005	24,726	7.8	8.2	24,324	11.1	21,339	2.9	24,996	62.3
	2004	28,209	7.6	8.6	27,611	11.4	24,125	3.0	28,189	62.2
	2003	0	*	*	0	*	0	*	0	*
	2002	23,758	7.4	9.2	23,341	10.7	20,350	3.1	23,977	59.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

